

[Cat. No.] Please refer to the **Ordering Information**

Introduction

AccuPower® RocketScript™ RT-PCR PreMix utilizes one-step RT-PCR performing cDNA synthesis and PCR in a single step, in a single tube. It can effectively synthesize cDNA with complex secondary RNA structures by using RocketScript™ Reverse Transcriptase (RocketScript™ RTase) developed by BIONEER. RocketScript™ RTase originated and engineered from M-MLV reverse transcriptase can provide increased thermal stability resulting in effective full length first strand cDNA synthesis. It can be used at various temperatures (42-70°C) allowing increased specificity, higher yields of cDNA, and more full-length product than other reverse transcriptase. This product contains vacuum-dried components including RocketScript™ RTase, reaction buffer, DTT, dNTPs, RNase inhibitor, and Top DNA Polymerase. It simplifies preparation of reverse transcription reaction mixture and PCR mixture by adding template RNA and primers without any extra process. After the reaction, samples can be applied directly on agarose gel for analysis due to included tracking dye.




Figure 1. Schematic diagram of 5' UTR of a gene, with complex secondary structure, at three different temperatures.

RocketScript™ Reverse Transcriptase is fully active at 70°C, so it can synthesize complete gene sequences that M-MLV and other reverse transcriptase can not synthesize.

Applications

- First-strand synthesis of cDNA from RNA molecules (RT)
- RT-PCR
- cDNA library construction
- Gene expression analysis

Features & Benefits

- Long range RT-PCR: Increased fidelity and accuracy enables one-step RT-PCR of full-length size of target up to 6 kb.
- Sensitivity: Excellent sensitivity and efficiency of cDNA synthesis even with small amount of RNA (10 pg-5 µg).
- Thermostable activity: RocketScript™ RTase has excellent thermal resistance even at 70°C, so it is suitable for cDNA synthesis of secondary RNA structures. The cDNA synthesis temperatures can be set at various temperatures (42-70°C) according to user's experimental conditions, allowing effective cDNA synthesis.
- Reproducibility: Mass production under ISO 9001 quality system allows minimized deviation between lots and reproducible results in replicated tests performed under same conditions and variation.
- Ease-of-use: Reactants are individually packaged in each of the PCR tubes, it allows any user simply perform cDNA synthesis and PCR by adding template RNA and primers. In addition, tracking dye and sedimentation agents are included in products for electrophoresis.

Composition

Composition	Concentration
RocketScript™ Reverse Transcriptase	200 U
5X Reaction buffer	1X
DTT	0.25 mM
dNTPs (dATP, dCTP, dGTP, dTTP)	Each 250 µM
RNase inhibitor	1 U
Top DNA Polymerase	1 U
Stabilizer and tracking dye	1X

Specifications

Top DNA Polymerase	
5' to 3' exonuclease activity	No
3' to 5' exonuclease activity	No
3'-A overhang	Yes
Fragment size	Up to 6 kb

Storage

Store at -20°C. If stored in the recommended temperature, this product will be stable until the expiration date printed out on the label.

Online Resources



Korean



English

Visit our **product page** for additional information and protocols


Ordering Information

Description	Cat. No.
0.2 ml thin-wall 8-tube strips with attached cap	20 µl/rxn K-2501 50 µl/rxn K-2503 20 µl/rxn K-2502 50 µl/rxn K-2504




Notice

BIONEER corporation reserves the right to make corrections, modifications, improvements and other changes to its products, services, specifications or product descriptions at any time without notice.

Explanation of Symbols



Experimental Procedures

Steps		Procedure Details																												
1	 Preparation of reaction mixture	<p>1. Add template RNA, primers and nuclease-free water into <i>AccuPower® RocketScript™</i> RT-PCR PreMix tubes to make a total volume of 20 µl or 50 µl. Do not include the dried pellet.</p> <ul style="list-style-type: none"> Amount of template RNA and primers <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th colspan="2">Components</th> <th>20 µl reaction</th> <th>50 µl reaction</th> </tr> </thead> <tbody> <tr> <td rowspan="2">Template RNA</td> <td>Total RNA</td> <td>0.01-5 µg</td> <td>0.01-5 µg</td> </tr> <tr> <td>Poly(A) RNA</td> <td>0.01-5 µg</td> <td>0.01-5 µg</td> </tr> <tr> <td>Primers</td> <td>Gene specific primer</td> <td>10-30 pmol</td> <td>10-30 pmol</td> </tr> </tbody> </table> <p>2. Dissolve the vacuum-dried pellet by pipetting or vortexing, and briefly spin down.</p>	Components		20 µl reaction	50 µl reaction	Template RNA	Total RNA	0.01-5 µg	0.01-5 µg	Poly(A) RNA	0.01-5 µg	0.01-5 µg	Primers	Gene specific primer	10-30 pmol	10-30 pmol													
Components		20 µl reaction	50 µl reaction																											
Template RNA	Total RNA	0.01-5 µg	0.01-5 µg																											
	Poly(A) RNA	0.01-5 µg	0.01-5 µg																											
Primers	Gene specific primer	10-30 pmol	10-30 pmol																											
2	 RT-PCR	<p>3. Perform the reaction under the following conditions.</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th>Step</th> <th>Temperature</th> <th>Time</th> <th>Cycles</th> </tr> </thead> <tbody> <tr> <td>cDNA synthesis</td> <td>42-70°C</td> <td>10-60 min</td> <td>1 cycle</td> </tr> <tr> <td>Pre-denaturation</td> <td>95°C</td> <td>5 min</td> <td>1 cycle</td> </tr> <tr> <td>Denaturation</td> <td>95°C</td> <td>10-30 sec</td> <td></td> </tr> <tr> <td>Annealing</td> <td>50-65°C</td> <td>10-30 sec</td> <td>30 cycles</td> </tr> <tr> <td>Extension</td> <td>72°C</td> <td>1 kb/min</td> <td></td> </tr> <tr> <td>Final extension</td> <td>72°C</td> <td>5 min</td> <td>1 cycle</td> </tr> </tbody> </table>	Step	Temperature	Time	Cycles	cDNA synthesis	42-70°C	10-60 min	1 cycle	Pre-denaturation	95°C	5 min	1 cycle	Denaturation	95°C	10-30 sec		Annealing	50-65°C	10-30 sec	30 cycles	Extension	72°C	1 kb/min		Final extension	72°C	5 min	1 cycle
Step	Temperature	Time	Cycles																											
cDNA synthesis	42-70°C	10-60 min	1 cycle																											
Pre-denaturation	95°C	5 min	1 cycle																											
Denaturation	95°C	10-30 sec																												
Annealing	50-65°C	10-30 sec	30 cycles																											
Extension	72°C	1 kb/min																												
Final extension	72°C	5 min	1 cycle																											
3	 Analyze with gel electrophoresis	<p>4. After the reaction, maintain the reaction mixture at 4°C. The samples can be stored at -20°C until use.</p> <p>5. Load samples on agarose gel without adding a loading-dye mixture, and perform gel electrophoresis for analysis.</p>																												