

[Cat. No.] TA-1019-1

## Introduction

Bioneer AccuNanoBead Thiol Magnetic NanoBeads are uniform, silica-based paramagnetic beads coated with high density thiol functional groups on the surface. The beads are used to reversible couple thiol-containing ligands. Thiol Magnetic Beads are most suitable for conjugation of large proteins.

## Features & Benefits

- Recommended coupling conditions: pH 4–8, 4°C to 25°C, 3–16 h.
- Specific isolation of cysteine proteins/peptides
- Stable covalent bond with minimal ligand leakage
- Produces reusable immunoaffinity matrices
- Low nonspecific binding
- Applications: Cell sorting, Immunoprecipitation; Purification for Antibodies, Proteins/Peptides, DNA/RNA

## Components

Components	Amount
AccuNanoBead™ Thiol Magnetic NanoBeads	0.5 g/ 25ml in 20% Ethanol

\* **Note:** For research use only. Not for use in diagnostic or therapeutic procedures.

## Materials to be Prepared by User

Magnetic Separator	
Coupling Buffer	0.1 M sodium phosphate, pH 7.0 , 5mM EDTA
L-Cysteine•HCl	
TCEP(tris(2-carboxyethyl)phosphine)	
Washing Buffer	1 M NaCl, 0.05% NaN3

\* **Note:** Buffer could be changed depending on user's needs.

## Specifications

AccuNanoBead™ Thiol Magnetic NanoBeads	
Composition	Thiol Magnetic NanoBeads
Binding capacity	≥ 400 nmol/g-beads
Size	Average 400 nm
Concentration	0.5 g(Solid)

## Storage

Store at room temperature.

This product can be stable for 3 years at room temperature (25°C).

## Expired date

Indicated on the label.

## Precautions

- Do not vigorously vortex AccuNanoBead™ Thiol Magnetic NanoBeads
- An exact protocol may need to be optimized by the user

## Online Resources



Korean



English

Visit our product page for additional information and protocols


## Ordering Information

Description	Cat. No.
AccuNanoBead™ Thiol Magnetic NanoBeads	TA-1019-1

## Notice

BIONEER corporation reserves the right to make corrections, modifications, improvements and other changes to its products, services, specifications or product descriptions at any time without notice.

## Explanation of Symbols



**Experimental Procedures (The protocols are scalable and can be optimized)**

Steps		Procedure Details
1	<b>Coupling</b>	<ol style="list-style-type: none"> <li>1. Add the protein to the coupling buffer.</li> <li>2. Put the magnetic beads in the tube, add coupling buffer and disperse the beads using an ultrasonicator.</li> <li>3. Add the protein sample prepared in step 1 to the tube with dispersed magnetic beads.</li> <li>4. Rotate the dispersed bead-proteins in the tube on a rotator for 30 minutes at room temperature.</li> <li>5. Remove the supernatant by holding the magnet close to the tube, and reload the buffer.</li> <li>6. Wash steps 4 through 5 with coupling buffer 3 times.</li> <li>7. Disperse the beads in 1ml coupling buffer containing L-cysteine•HCl, and rotate on a rotator for 30 minutes at room temperature.</li> <li>8. Place the magnet close to the tube to remove the supernatant and wash 4 times with Washing buffer.</li> <li>9. Disperse beads in PBS buffer containing 0.05% sodium azide and store at 4°C.</li> </ol>